



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Reply: Expression of the proteolysis-inducing factor core-peptide mRNA is upregulated in both tumour and adjacent normal tissue in gastroesophageal malignancy

Citation for published version:

Deans, DAC, Wigmore, SJ, Gilmour, H, Tisdale, MJ, Fearon, KCH & Ross, JA 2008, 'Reply: Expression of the proteolysis-inducing factor core-peptide mRNA is upregulated in both tumour and adjacent normal tissue in gastroesophageal malignancy', *British Journal of Cancer*, vol. 98, no. 1, pp. 243-243.
<https://doi.org/10.1038/sj.bjc.6604117>

Digital Object Identifier (DOI):

[10.1038/sj.bjc.6604117](https://doi.org/10.1038/sj.bjc.6604117)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

British Journal of Cancer

Publisher Rights Statement:

Available under Open Access

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Letter to the Editor

Reply: Expression of the proteolysis-inducing factor core-peptide mRNA is upregulated in both tumour and adjacent normal tissue in gastrooesophageal malignancy

DAC Deans¹, SJ Wigmore¹, H Gilmour¹, MJ Tisdale¹, KCH Fearon¹ and JA Ross^{*,1}¹Tissue Injury and Repair Group, Clinical and Surgical Sciences, University of Edinburgh, The Chancellor's Building, 49 Little France Crescent, Edinburgh EH16 4SB, UK

British Journal of Cancer (2008) 98, 243. doi:10.1038/sj.bjc.6604117 www.bjcancer.com

Published online 18 December 2007

© 2008 Cancer Research UK

Sir,

Extrapolating from our findings that mRNA for the dermcidin gene (Schitteck *et al*, 2001), which encodes the proteolysis-inducing factor (PIF) core peptide (PIF-CP), is present in both gastrooesophageal tumour and adjacent non-tumour tissue, EA Rapoport suggests that PIF may be generally involved in muscle breakdown via production of PIF in tissues requiring amino acids for tissue regeneration, for example, in liver regeneration. There are, however, several caveats, which need to be applied to this suggestion. The first is that it is only the glycosylated form of the PIF-CP that has been shown to induce muscle breakdown (Todorov *et al*, 1996), and this form is difficult to identify in tissues (Wieland *et al*, 2007). Glycosylated PIF has also been shown to have pro-inflammatory effects on hepatocytes (Watchorn *et al*, 2001) and other cell types. In our study (Deans *et al*, 2006), we made no claim that glycosylated PIF was present in gastro-

oesophageal tumours or adjacent tissues. In addition, the PIF-CP is homologous to a neural survival peptide (YP-30) (Cunningham *et al*, 2002), and expression of the gene encoding the PIF-CP has been shown to provide a small survival and proliferative advantage to other cell types including tumours (Lowrie *et al*, 2006, and Stewart *et al*, 2007). Furthermore, evidence suggests that the YP-30/PIF-CP peptide may have a role in development, acting as a maternal blood-borne factor, which promotes survival of the developing thalamus (Landgraf *et al*, 2005). In a recent development, there is now evidence that peptide products of the dermcidin gene may participate in the regulation of placental function by means of modulating proteolytic cascades on the trophoblastic surface (Motoyama *et al*, 2007) and having proteolytic activity. Therefore, while EA Rapoport's suggestion may be unlikely in the light of present evidence, there is still much to be discovered about this fascinating gene and its products.

REFERENCES

- Cunningham TJ, Jing H, Akerblom I, Morgan R, Fisher TS, Neveu M (2002) Identification of the human cDNA for new survival/evasion peptide (DSEP): studies *in vitro* and *in vivo* of overexpression by neural cells. *Exp Neurol* 177: 32–39
- Deans DA, Wigmore SJ, Gilmour H, Tisdale MJ, Fearon KC, Ross JA (2006) Expression of the proteolysis-inducing factor core peptide mRNA is upregulated in both tumour and adjacent normal tissue in gastrooesophageal malignancy. *Br J Cancer* 94: 731–736
- Landgraf P, Sieg F, Wahle P, Meyer G, Kreutz MR, Pape HC (2005) A maternal blood-borne factor promotes survival of the developing thalamus. *FASEB J* 19: 225–227
- Lowrie AG, Wigmore SJ, Wright DJ, Waddell ID, Ross JA (2006) Dermcidin expression in hepatic cells improves survival without N-glycosylation, but requires asparagine residues. *Br J Cancer* 94: 1663–1671
- Motoyama J-PL, Kim-Motoyama H, Kim P, Nakagama H, Miyagawa K, Suzuki K (2007) Identification of dermcidin in human gestational tissue and characterization of its proteolytic activity. *Biochem Biophys Res Commun* 357: 828–833
- Schitteck B, Hipfel R, Sauer B, Bauer J, Kalbacher H, Stevanovic S, Schirle M, Schroeder K, Blin N, Meier F, Rassner G, Garbe C (2001) Dermcidin: a novel human antibiotic peptide secreted by sweat glands. *Nat Immunol* 2: 1133–1137
- Stewart GD, Lowrie AG, Riddick AC, Fearon KC, Habib FK, Ross JA (2007) Dermcidin expression confers a survival advantage in prostate cancer cells subjected to oxidative stress or hypoxia. *Prostate* 67: 1308–1317
- Todorov PT, McDevitt TM, Cariuk P, Coles B, Deacon M, Tisdale MJ (1996) Induction of muscle protein degradation and weight loss by a tumor product. *Cancer Res* 56: 1256–1261
- Watchorn TM, Waddell ID, Dowidar N, Ross JA (2001) Proteolysis inducing factor regulates hepatic gene expression via the transcription factors NFκB and STAT3. *FASEB J* 15: 562–564
- Wieland BM, Stewart GD, Skipworth RJE, Sangster K, Fearon KCH, Ross JA, Reiman TJ, Easaw J, Mourtzakis M, Kumar V, Pak BJ, Calder K, Filippatos G, Kremastinos DT, Palcic M, Baracos VE (2007) Is there a human homolog to the murine proteolysis inducing factor? *Clin Cancer Res* 13: 4984–4992

*Correspondence: Dr JA Ross; E-mail: j.a.ross@ed.ac.uk

Published online 18 December 2007